

ANALYTICA CHIMICA ACTA

An international journal devoted to all branches of analytical chemistry

(Full texts are incorporated in CJESEVIER, a file in the Chemical Journals Online database available on STN International; Abstracted, indexed in: Aluminium Abstracts; Anal. Abstr.; Biol. Abstr.; BIOSIS; Chem. Abstr.; Curr. Contents Phys. Chem. Earth Sci.; Engineered Materials Abstracts; Excerpta Medica; Index Med.; Life Sci.; Mass Spectrum. Bull.; Material Business Alerts; Metals Abstracts; Sci. Citation Index)

VOL. 409 NOS. 1-2

CONTENTS

March 23rd, 2000

Letter

In vivo or in vitro microdialysis?

- L. Gorton (Lund, Sweden) and N. Torto (Gaborone, Botswana) 1

Sequential injection

Development of a sequential injection system in the capillary format for determinations of the IpaC protein

- D.M. Spence, E.D. Knoll, B.T. Ruotolo and J. Bjerregaard (Saint Louis, MO, USA) 3

Kinetic-turbidimetric determination of phytic acid by sequential injection analysis

- J.G. March, B.M. Simonet and F. Grases (Palma de Mallorca, Spain) 9

Chromatography and electrophoresis

Ion chromatography using a charged complex anion-exchange group

- N. Hirayama, W. Umehara, H. Makizawa and T. Honjo (Kanazawa, Japan) 17

Sulfide measurements by flow injection analysis and ion chromatography with electrochemical detection

- I.G. Casella, M.R. Guascito (Potenza, Italy) and E. Desimoni (Milan, Italy) 27

Separation and determination of niobium(V) and tantalum(V) as 2-(5-bromo-2-pyridylazo)-5-[N-propyl-N-(3-sulfopropyl)amino]phenol citrate ternary complexes in geological samples using ion interaction high-performance liquid chromatography

- N. Vachirapatama, P. Doble and P.R. Haddad (Tasmania, Australia) 35

An HPLC detection scheme for underivatized amino acids based on tryptophan fluorescence recovery

- M. Yang and S.A. Tomellini (Durham, NH, USA) 45

Capillary electrophoretic determination of coordination equilibrium constants of carboxylate ions with copper(II) complexed with several bis(1,10-phenanthroline) derivatives and bis(bipyridine)


- T. Yokoyama, H. Tsuji and M. Zenki (Okayama, Japan) 55

Chemiluminescence

Chemiluminescence behavior of the decomposition of hydrogen peroxide catalyzed by copper(II)-amino acid complexes and its application to the determination of tryptophan and phenylalanine

- S. Hanaoka, J.-M. Lin and M. Yamada (Tokyo, Japan) 65

(Continued overleaf)



<http://chemweb.com/gateways/elsevier.html>

CONTENTS
Direct

This journal is part of **ContentsDirect**, the **free** alerting service which sends tables of contents by e-mail for Elsevier Science books and journals. You can register for **ContentsDirect** online at: www.elsevier.nl/locate/contentsdirect

| | |
|--|-----|
| Studies of micelle and trace non-polar organic solvent on a new chemiluminescence system and its application to flow injection analysis G.-F. Zhang and H.-Y. Chen (Nanjing, China) | 75 |
| Coulometric microflow titrations with chemiluminescent and amperometric equivalence point detection. Bromimetric titration of low concentrations of hydrazine and ammonium Z.K. He, B. Fuhrmann and U. Spohn (Halle, Germany) | 83 |
| <i>Electroanalytical chemistry</i> | |
| Determination of lead (II) and cadmium (II) in hard and soft wheat by derivative potentiometric stripping analysis F. Lo Coco (Udine, Italy), P. Monotti (Perugia, Italy), V. Fiecchi (Venice, Italy) and L. Ceccon (Udine, Italy) | 93 |
| Determination of liposomes drug retention capacity using differential pulse polarography: the case of chlorothiazide C.G. Kontoyannis and D. Douroumis (Rio-Patras, Greece) | 99 |
| Voltammetric behaviour and determination of the anthracycline antitumor drug nogalamycin M.S. Ibrahim (Al-Ain, United Arab Emirates) | 105 |
| Flow electrochemical determination of ascorbic acid in real samples using a glassy carbon electrode modified with a cellulose acetate film bearing 2,6-dichlorophenolindophenol A.B. Florou, M.I. Prodromidis, M.I. Karayannis and S.M. Tzouwara-Karayanni (Ioannina, Greece) | 113 |
| Pseudo-equilibrium approach to the design and use of enzyme-based amperometric biosensors evaluated using a sensor for hydrogen peroxide W. Chen and H.L. Pardue (West Lafayette, IN, USA) | 123 |
| [60]Fullerene-mediated amperometric biosensors V.G. Gavalas and N.A. Chaniotakis (Iraklion, Greece) | 131 |
| Response of nanosized cobalt oxide electrodes as pH sensors L. Qingwen, L. Guoan (Beijing, PR China) and S. Youqin (Zhengzhou, PR China) | 137 |
| Gas-gap sensors for the determination of nitrogen oxides and nitrites O.V. Yagodina, E.B. Nikolskaya and N.B. Shor (St. Petersburg, Russia) | 143 |
| <i>Immunoassay</i> | |
| Flow injection electrochemical enzyme immunoassay based on the use of gold bands E.M. Abad-Villar, M.T. Fernández-Abedul and A. Costa-García (Asturias, Spain) | 149 |
| <i>Chemometrics and experimental design</i> | |
| Application of two- and three-way chemometric methods in the study of acetylsalicylic acid and ascorbic acid mixtures using ultraviolet spectrophotometry M.M. Sena, J.C.B. Fernandes, L. Rover Jr., R.J. Poppi and L.T. Kubota (Campinas, Brazil) | 159 |
| Trimethyltin chloride determination by hydride generation — gas phase molecular absorption spectrometry. A sequential application of experimental designs J. Sanz-Asensio, M.T. Martínez-Soria, M. Plaza-Medina and M. Pérez-Clavijo (La Rioja, Spain) | 171 |
| Exhaustive set of non-isomorphic sub-graphs and its application to chemical structure elucidation using a IR spectroscopy database V.N. Piottukh-Peletsky, I.K. Korobeinicheva, T.F. Bogdanova, S.G. Molodtsov and B.G. Derendyaev (Novosibirsk, Russia) | 181 |
| <i>Molecular emission cavity analysis</i> | |
| New cavity design suitable for monitoring gaseous samples by molecular emission cavity analysis N. Maleki, A. Safavi and Z. Ramezani (Shiraz, Iran) | 197 |
| <i>Thermogravimetry – mass spectrometry</i> | |
| Quantitative thermogravimetric-mass spectrometric analysis for monitoring the effects of fire retardants on cellulose pyrolysis M. Statheropoulos and S.A. Kyriakou (Athens, Greece) | 203 |
| A shipboard method for arsenic speciation using semi-automated hydride generation atomic fluorescence spectroscopy A.M. Featherstone, P.R. Boulton, B.V. O'Grady and E.C.V. Butler (Tasmania, Australia) | 215 |
| A flow system exploiting multicommutation for speciation of inorganic nitrogen in waters F.R.P. Rocha (Piracicaba SP, Brazil and São Carlos, Brazil) and B.F. Reis (Piracicaba SP, Brazil) | 227 |
| Determination of triphenyltin in sea-water by excitation–emission matrix fluorescence and multivariate curve resolution J. Saurina, C. Leal, R. Compañó, M. Granados, R. Tauler and M.D. Prat (Barcelona, Spain) | 237 |
| Electrochemical and conformational studies of microcystin–LR F. Yan, M. Ozsoz and O.A. Sadik (Binghamton, NY, USA) | 247 |
| Isolation and identification of unsaturated fatty acid methyl esters from marine micro-algae C. Viron (Orléans, France and St. Jean de Braye, France), A. Saunois, P. André (St. Jean de Braye, France), B. Perly (Gif-sur-Yvette, France) and M. Lafosse (Orléans, France) | 257 |
| The use of tungsten–rhodium permanent chemical modifier for cadmium determination in decomposed samples of biological materials and sediments by electrothermal atomic absorption spectrometry É.C. Lima, F. Barbosa Jr. and F.J. Krug (Piracicaba, Brazil) | 267 |
| <i>Kinetic methods</i> | |
| Kinetic spectrophotometric determination of V(IV) in the presence of V(V) by the H-point standard addition method A. Safavi, H. Abdollahi, F. Sedaghatpour and S. Zeinali (Shiraz, Iran) | 275 |

| | |
|---|-----|
| Highly selective and sensitive kinetic spectrophotometric determination of vanadium(IV) in the presence of vanadium(V) A. Safavi, M.R. Hormozi Nezhad (Shiraz, Iran) and E. Shams (Zanjan, Iran) | 283 |
| <i>Author Index</i> | 291 |
| <i>Calendar of forthcoming meetings</i> | N1 |

